CUMULATIVE AUTHOR INDEX
Annals of the Entomological Society of America
Volume 101, 2008

A
Abraham, S., 639
Acaron, A., 964
Aiello, A., 689
Albuquerque, G. S., 314
Allison, C. J., 1162
Alyokhin, A. V., 1078
Andersen, P. C., 1033
Anfora, G., 1003
Appel, A. G., 397
Armstrong, J. S., 1112
Ascher, J. S., 140, 1067

B
Backus, E. A., 159
Bagwell, G. J., 235
Baker, M. B., 371
Baldissera, V. D., 619
Barra, J.-A., 501
Bartlett-Healy, K., 95
Baselga, A., 402
Becerra-Chiron, I. M., 113
Bechinski, E. J., 351
Beitz, D. C., 430
Benito, J.C.S., 683
Bento, J.M.S., 551
Bernal, J. S., 113
Bidau, C. J., 802
Birand, A. C., 30
Bochkov, A. V., 289
Booth, W., 763
Borovec, R., 7
Boyd, E. A., 253
Brailovsky, H., 307
Brandt, S. L., 159
Breed, M. D., 1145
Broce, A. B., 650
Brown, B. V., 713
Brown, R. L., 899
Buhl, P. N., 297
Burns, J. A., 1026
Burns, J. M., 794
Bustamante, D. M., 867

C
Calcaterra, L. A., 539
Calvo, D., 881
Cambra T., R. A., 689
Campbell, D. R., 70
Campos, L.A.O., 915
Cardé, R. T., 215
Carlton, C. E., 13
Carrasco, J. V., 181
Castrejón Gómez, V. R., 181
Caterino, M. S., 939
Chacón-Torres, N. M., 113
Chatzimanolis, S., 939
Chen, X.-X., 189
Cheng, C.-W., 950
Cheng, I.-C., 950
Christiansen, K., 501, 989
Cladera, J. L., 627
Coats, J. R., 430
Coderre, D., 860
Čokl, A., 235, 1169
Cole, B. J., 272
Cole, J. A., 815
Contreras-Ramos, A., 703, 808
Conway, M., 964
Cordero, C., 786
Coudron, T. A., 159, 613
Crans, W., 95
Cristofaro, M., 7
Crook, D. J., 1094, 1103
Cuda, J. P., 604
Cueva del Castillo, R., 260

D
Dash, S. T., 1056
Dastur, S. R., 371
Davis, D. R., 689
De Freitas Fialho, J., 779
Delgado, A., 539
Delgado, L., 53
Demite, P. R., 804
Deyrup, M., 1179
Dias, C. M., 1145
Diaz, R., 604
Dickenson, R. M., 229
Dixon, P. M., 341
Djam, S., 930
Doberlet, M. V., 1169
Dormont, L., 845
Dunkley, J. P., 867
Duso, C., 297

E
Edwards, R., 272
Ekesi, S., 331
Emmert, C. J., 1033
Enstrom, P. C., 1154
Er, A., 593
Ergin, E., 593
Escobar, L. A., 867
Evans, C. S., 664
Evans, G., 525

F
Faccio, P., 627
Favret, C., 833
Felgenhauer, B. E., 1026
Feres, R.J.F., 804
Fernandes, P. M., 551
Ferris, V. R., 204
Fetridge, E. D., 1067
Fielding, D. J., 439
Finlayson, C. J., 1078
Fiúza, L. M., 619
Fordyce, J. A., 172, 794
Forister, M. L., 794
Fox, A., 604
Frank, J. H., 1033
Fraser, S., 418
Frizzas, M. R., 779
Fuchikawa, T., 1121
Funderburk, J. E., 769
Furniss, M. M., 1010

G
Galicia, I., 786
Garczynski, S. F., 887
Gaugler, R., 95
Gautam, B. K., 1088
Geertsema, H., 197
Gende, P., 930
Gereszek, L. J., 430
Gillette, N. E., 845
Girling, R. D., 215
Gómez-Gómez, B., 128
Gordon, I., 331
Greenberg, L., 1162
Grossman, J. D., 266
Grzywacz, B., 837
Guédot, C., 675
Gullan, P. J., 779
Gültekin, L., 7

H
Habibi, J., 159
Halbert, S. E., 20
Hall, D. G., 585
Hance, T., 598
Harmer, A.M.T., 664
Harris, J. W., 1137
Haruyama, N., 971
He, Y., 1017
Head, G. P., 639
Hebbar, P. K., 930
Henderson, G., 1088
Henry, T. J., 20
Heraty, J. M., 491
Hernández, M. C., 1041
Heyward, J., 272
Hoddle, M. S., 253, 491
Holbrook, C. T., 272

Holm, L., 272
Holzenthal, R. W., 465
Honek, A., 533
Hooper-Bùi, L. M., 1056
Horng, S.-B., 950
Horton, D. R., 887
Howard, D. J., 30
Hoy, M. A., 1050
Huang, F., 189
Huesing, J. E., 159
Hunter, R. K., 1026
Hurvois, J.-P., 1162

I
Iannuzzi, R., 79
Ioriatti, C., 1003
Ishizaki, M., 558
Itoh, T. Q., 1121

J
Jaffe, B. D., 371
Jay-Robert, P., 58
Jiang, H.-B., 1017
Jiménez, M., 993
Jones, R. W., 128
Jordana, R., 501
Juaréz, M. L., 639
Juliano, S. A., 1150

K
Kabashima, J. N., 1162
Kaminski, L. A., 906
Kegley, S. J., 1010
Kerr, L. M., 1094, 1103
Kesavaraju, B., 1150
King, B. H., 229
Klotz, J. H., 1162
Knisley, C. B., 277
Komazaki, S., 565
Kowalczyk, M., 837

L
Labandeira, C. C., 79
Labrie, G., 860
Lambert, S., 930
Landolt, P. J., 675
Landry, K. M., 1078
Langellotto, G. A., 140, 1067
Lanzavecchia, S. B., 627
Laup, S., 930
Lazarowitz, S. R., 1026
Leathers, J. W., 682
Lebreton, J.-D., 58
Lee, C.-F., 722
Legaspi, B. C., 378
Legaspi, J. C., 378
Lemon, M., 664
Leschen, R.A.B., 13
Li, C.-L., 474
Li, H.-F., 456
Li, J., 613
Li, S., 204
Li, W., 613
Liu, X., 613
Lofego, A. C., 804
Lognay, G. C., 598
Lonsdale, O., 327
Lu, Z., 613
Lucas, É., 860
Lucas, L. K., 172
Lucchi, A., 1003
Luckhart, S., 824
Lumaret, J.-P., 58

M
Ma, Z., 351
Machado, V., 619
Maisin, N., 930
Mankin, R. W., 664
Mano, H., 449
Marcos-García, M. Á., 384
Marshall, J. L., 30
Marshall, S. A., 327
Martí, D. A., 802
Martins De Oliveira, C., 779
Marty, N. P., 627
Mastro, V. C., 1094, 1103
Masumoto, K., 474
Matsumoto, A., 1121
Matsuyama, T., 1121
Matteson, K. C., 140
Mazzoni, V., 1003
McCabe, T. L., 40
McElfresh, J. S., 215
McKay, T., 650
Meagher, R. L., 1112
Mendez, P. K., 573
Menzel, T. O., 119
Messing, R. H., 656
Millar, J. G., 215, 235, 1169
Miller, D., 525
Miyashita, T., 392
Miyatake, T., 1121
Mizell, R. F., 1033
Mochizuki, A., 971
Molina, J. M., 881
Moore, M. K., 1026
Mound, L. A., 491
Moya-Raygoza, G., 113
Muratori, F., 598
Murúa, M. G., 639

N
Nagoshi, R. N., 619, 1112
Naka, H., 971
Nardi, C., 551
Nebeker, T. E., 119
Neves, C. A., 915
Nice, C. C., 172, 794
Nielsen, C., 833
Nigg, H. N., 418
Nomura, M., 971
Northfield, T. D., 769
Novoa, F., 402

O
O'Brien, C. W., 128
Ochi, T., 474
OConnor, B. M., 289
Ogden, T. H., 1
Ogol, C.K.P.O., 331
Ohta, Y., 1121
Oliveira, J. V., 619
O'Neal, M. E., 341
Oseto, C. Y., 204
Overholt, W. A., 604

P
Paine, T. D., 1162
Paini, D. R., 769
Palacios-Vargas, J. G., 683
Palomera-Avalos, V., 113
Pan, W., 613
Paretas-Martínez, J., 993
Percy, D. M., 253
Peredo, L. C., 307
Pfannenstiel, R. S., 763
Pfeiffer, D. G., 824
Phillips, T. W., 955
Pineda, A., 384
Pitts, J. P., 514
Pogue, M. G., 743
Ponce-Wainer, J. X., 260
Pratt, P. D., 604
Prieto, S., 639
Proud, D. N., 1026
Pujade-Villar, J., 993
Purung, H. b., 930

Q
Quintero A., D., 689

R
Reithel, J. S., 70
Reitz, S. R., 769
Remis, M. I., 922
Resh, V. H., 573
Reynoso-Velasco, D., 703
Rhainds, M., 872
Robertson, D. R., 465
Ronnau, M., 915
Roques, A., 845
Rosetti, N., 922
Rosmana, A., 930
Rost-Roszkowska, M. M., 151
Rota, J., 40
Roux-Morabito, G., 845
Rugman-Jones, P. F., 491
Ruiz-Montoya, L., 128
Rung, A., 525
Rust, M. K., 1162
Rwomushana, I., 331

S
Sadof, C., 872
Sambaraju, K. R., 955

Sanada-Morimura, S., 1121
Sánchez, V., 786
Sartori, M., 1
Saska, P., 533
Schaefer, C. W., 463
Schal, C., 763
Scheffer, S. J., 525, 930
Schmidt, N. P., 341
Schoenick, C. A., 1154
Schuh, R. T., 20
Serrão, J. E., 915
Shapiro, A. M., 794
Shapiro, L. H., 930
Shi, M., 189
Silvie, P., 1112
Simmons, A. M., 378
Simmons, R. B., 743
Simonsen, T. J., 899
Skarżyński, D., 989
Smith, L., 7
Smith, P. T., 713
Smith, R. J., 266
Snoddy, E. T., 397
Song, D.-X., 1131
Soto-Adames, F. N., 501
Sperling, F.A.H., 845, 899
Stein, J., 845
Stimac, J. L., 1033
Stireman, J. O., III, 362
Stoffolano, J. G., Jr, 964
Stouthamer, R., 491
Su, N.-Y., 456
Suasa-Ard, W., 301
Sulistyowati, E., 930
Sun, L., 204
Switzer, P. V., 1154

T

Takagi, S., 392
Tang, P.-A., 1017
Tanimura, T., 1121
Tatsuta, H., 837
Tauber, C. A., 314
Tauber, M. J., 314
Taylor, P. W., 664
Tibbetts, J. A., 1026
Timm, A. E., 197
Toda, S., 565
Tokuda, M., 301
Toquenaga, Y., 449
Townsend, V. R., Jr, 1026
Traylor, T., 30
Tsutsui, N. D., 539
Tu, Z., 824
Tüven, A., 593

U

Uçkan, F., 593
Unruh, T. R., 887

V

Van Dam, A. R., 411
Van Dam, M. H., 411
Vargo, E. L., 763
Vega, F. E., 930
Vera, M. T., 639
Vilardi, J. C., 627, 922
Villa, J. D., 867
Vogler, A. P., 277

W

Wagner, D. L., 40
Wagner, R. M., 159
Wang, C.-C., 474
Wang, J.-J., 1017
Wang, X.-G., 656
Wang, Z.-Y., 1017
Warchałowska-śliwa, E., 837
Warnich, L., 197
Watanabe, T., 558
Weirauch, C., 20
Wenninger, E. J., 585
Whiting, M. F., 1
Wiernasz, D. C., 272
Willett, L. S., 887
Willink, E., 639
Wilson, J. S., 514
Wolak, M. E., 801
Wong, T., 371
Woodcock, M. R., 277
Wright, M. K., 159
Wunder, M., 619

Y

Yang, P.-S., 474
Yang, R.-L., 950
Yano, S., 247
Yassin, A., 978
Yasuda, T., 558
Young, O. P., 104
Yu, R.-X., 189
Yu, T., 887
Yukawa, J., 301

Z

Zahn, D. K., 215
Zandomeni, R. O., 627
Zanuncio, J. C., 915
Zaspel, J. M., 1050
Zhang, J.-Y., 1131
Zhang, Q., 613
Zhang, X., 824
Zhou, K.-Y., 1131
Žunič, A., 1169

CUMULATIVE KEY WORD INDEX
Annals of the Entomological Society of America
Volume 101, 2008

A

Abies guatemalensis,
aphid, 833
conservation, 833
Guatemala, 833
abundance,
Atlantic forest, 402
Coleoptera, 402
conservation, 402
richness, 402
acoustic communication,
Pentatomidae, 235
sex-specific signal, 235
substrate-borne vibrational signal, 235
acoustic signal,
Homalodisca coagulata, 253
leafhopper, 253
sharpshooter, 253
substrate vibration, 253
activity,
foraging, 272
foraging effort, 272
foraging success, 272
Pogonomyrmex occidentalis, 272
adult survival,
B-chromosomes, 922
body-size related traits, 922
grasshoppers, 922
Aedes albopictus,
Corethrella appendiculata, 1150
predator–prey, 1150
size-dependent, 1150
AFLP,
dispersal, 197
gene flow, 197
Grapholita molesta, 197
population genetic structure, 197
Africa paleobiogeography,
COII barcoding, 978
cryptic speciation, 978
interspecific hybridization, 978
wing shape, 978
aggression,
competition strategy, 449
endophagous insect, 449
intraspecific competition, 449
pupal cell, 449
agricultural,
aphidophagous, 1078
biological control, 1078
conservation, 1078
invasive, 1078
agriculturally important species,
cryptic species, 743
mitochondrial DNA, 743
morphology, 743
phylogeny, 743
alary polymorphism,
below-ground insects, 551
brachypterous, 551
macropterous, 551
Alcaligenes,
blood-feeding moth, 1050
gut endosymbiont, 1050
Klebsiella, 1050
microbial associate, 1050
Allonemobius,
conspecific sperm precedence, 30
ground cricket, 30
hybrid zone, 30
hybridization, 30
amino acid,
Coptotermes formosanus, 1088
feeding deterrent, 1088
wood preservative, 1088
amphibian,
attraction, 95
Culex territans, 95
mosquito, 95
mosquito hearing, 95
amplification process,
collective dispersal, 247
solitary behavior, 247
Tetranychus urticae, 247
trail following, 247
Amuroclusia syn. nov.,
Clusia, 327
Clusiidae, 327
Melanoclusia gen. nov., 327
Paraclusia syn. nov., 327
androconial system,
caddisfly, 465
male genitalia, 465
Neotropics, 465
scales, 465
Annona cherimola,
antennal sensilla, 181
fruit borer, 181
Talponia batesi, 181
ant collection techniques,
extirpated species, 1056
faunal survey, 1056
pest species, 1056
state list, 1056
ant communities,
forest landscapes, 119
imported fire ants, 119
null models, 119
ant-organs,
balloon setae, 906
host plants, 906
myrmecoxeny, 906
scanning electron microscopy, 906
antennae,
contact chemoreceptors, 1094
electron microscopy, 1094
sensilla, 1094
antennal sensilla,
Annona cherimola, 181
fruit borer, 181
Talponia batesi, 181
ants,
interference competition, 1162
Linepithema, 1162
Solenopsis, 1162
venom, 1162
Apanteles galleriae,
gibberellic acid, 593
parasitoid, 593
risk assessment, 593
toxicity, 593
aphid,
Abies guatemalensis, 833
conservation, 833
Guatemala, 833
aphid epicuticule,
cuticular hydrocarbons, 598
exuviae, 598
Sitobion avenae, 598
wax esters, 598
aphidophagous,
agricultural, 1078
biological control, 1078
conservation, 1078
invasive, 1078
Apidae,
molecular systematics, 713
Neotropical, 713
parasitoid, 713
Apis mellifera,
recurrent event, 867
survival, 867
swarming, 867
Varroa destructor, 867
apple,
attraction, 675
codling moth, 675
host-finding, 675

pear, 675
Araeopteroninae,
Eublemminae, 40
Glenoides lenticuligera, 40
Hypenodinae, 40
Nigetia formosalis, 40
Archaeognatha,
complete mitochondrial DNA sequence, 1131
inversion, 1131
Pedetontus silvestrii, 1131
Asian citrus psyllid,
body color, 585
huanglongbing, 585
pigment, 585
reproductive state, 585
Asphonyliini,
Dimocarpomyia folicola, 301
Dimocarpus longan, 301
gall midges, 301
Thailand, 301
assemblages,
dung beetles, 58
southern alps, 58
spatial and temporal segregation, 58
Atlantic forest,
abundance, 402
Coleoptera, 402
conservation, 402
richness, 402
attraction, 95, 675
amphibian, 95
apple, 675
codling moth, 675
Culex territans, 95
host-finding, 675
mosquito, 95
mosquito hearing, 95
pear, 675
Australia,
Early Carboniferous, 79
orthopteroid, 79
pteridosperm, 79
time lag, 79

B

Bactrocera invadens,
fruit fly, 331
host plants, 331
host preference, 331
infestation index, 331
Bactrocera tryoni,
mating competitiveness, 664
sterile insect technique, 664
bagworm,
flightless Lepidoptera, 872
population regulation, 872
Thyridopteryx ephemeraeformis, 872
white pine, 872
balloon setae,
ant-organs, 906
host plants, 906
myrmecoxeny, 906
scanning electron microscopy, 906
B-chromosomes,
adult survival, 922
body-size related traits, 922
grasshoppers, 922
bee conservation,
biodiversity, 1067
parasitic bees, 1067
soil-nesting species, 1067
suburban development, 1067
bees, 915, 1137
brood, 1137
drone, 1137
hygiene, 1137
insect, 915
midgut, 915
peritrophic membrane, 915
varroa, 1137
behavioral sequences,
courtship, 215
harlequin bug, 215
Hemiptera, 215
Pentatomidae, 215
below-ground insects,
alary polymorphism, 551
brachypterous, 551
macropterous, 551
Bemisia tabaci,
biological control, 378
Delphastus catalinae, 378
predator, 378
survival, 378
bioacoustics,
Cicadidae, 815
cryptic species, 815
Tibicen, 815
biodiversity, 128, 1067
bee conservation, 1067
cloud forests, 128
conservation, 128
Curculionidae, 128
parasitic bees, 1067
soil-nesting species, 1067
suburban development, 1067
tropical montane forests, 128
bioecology,
cassava root scale, 779
Insecta, 779
Manihot esculenta, 779
biological control, 189, 378, 384, 539, 604, 1078
agricultural, 1078
aphidophagous, 1078
Bemisia tabaci, 378
Blissidae, 604
Braconidae, 189
climate, 539
conservation, 1078
degree-days, 604
Delphastus catalinae, 378
developmental rate, 604
greenhouses, 384
immature stages, 189
invasive, 1078
larval development, 189
modified Malaise trap, 384
parasitoid, 189
phenology, 539
Poaceae, 604
population dynamics, 384
predator, 378
Solenopsis invicta, 539
species relationships, 539
survival, 378
Syrphidae, 384
Blattella asahinai,
distribution map, 397
new distribution records, 397
peridomestic pest, 397
Blissidae,
biological control, 604
degree-days, 604
developmental rate, 604
Poaceae, 604
blood-feeding moth,
Alcaligenes, 1050
gut endosymbiont, 1050
Klebsiella, 1050
microbial associate, 1050
blow fly,
bubbling, 964
droplet production, 964
insect crop, 964
regurgitation, 964
body color,
Asian citrus psyllid, 585
huanglongbing, 585
pigment, 585
reproductive state, 585
body-size related traits,
adult survival, 922
B-chromosomes, 922
grasshoppers, 922
Bolboceratini,
Coleoptera, 474
Geotrupidae, 474
Scarabaeoidea, 474
brachypterous,
alary polymorphism, 551
below-ground insects, 551
macropterous, 551
Braconidae,
biological control, 189
immature stages, 189
larval development, 189
parasitoid, 189

Brazil,
carinulatus species group, 804
Galendromus (*Mugidromus*), 804
predatory mites, 804
South America, 804
brood,
bees, 1137
drone, 1137
hygiene, 1137
varroa, 1137
bubbling,
blow fly, 964
droplet production, 964
insect crop, 964
regurgitation, 964

C

C. carnea group,
courtship song, 971
cytochrome oxidase subunit I, 971
diagnostic marker, 971
mitochondrial polyphyly, 971
Cactoblastis cactorum,
dispersal, 899
introduction, 899
invasive species, 899
phylogeography, 899
caddisfly, 465, 573
androconial system, 465
case building, 573
life cycle, 573
male genitalia, 465
Neotropics, 465
phenology, 573
protandry, 573
scales, 465
calling song,
cricket, 260
mate choice, 260
predation, 260
Zapotitlán Salinas, 260
Callophrys,
Eueides, 786
Heliconius, 786
sexual coevolution, 786
spermatophore, 786
Callosobruchus maculatus,
dumping behavior, 950
egg-laying decision, 950
host range, 950
Calosoma sayi,
Carabidae, 104
food, 104
Spodoptera frugiperda, 104
starvation, 104
Carabidae,
Calosoma sayi, 104
food, 104
Spodoptera frugiperda, 104
starvation, 104
Caribbean,
ecology, 1026
Erythraeidae, 1026
Opiliones, 1026
parasitism, 1026
carinulatus species group,
Brazil, 804
Galendromus (*Mugidromus*), 804
predatory mites, 804
South America, 804
case building,
caddisfly, 573
life cycle, 573
phenology, 573
protandry, 573
cassava root scale,
bioecology, 779
Insecta, 779
Manihot esculenta, 779
Ceratitis capitata,
DNA molecular marker, 627
introduction sources, 627
mitochondrial haplotypes, 627
population dynamics, 627
chemical ecology,
chemical stimuli, 955
host selection behavior, 955
physical stimuli, 955
chemical stimuli,
chemical ecology, 955
host selection behavior, 955
physical stimuli, 955
chilling,
diapause, 439
Melanoplus, 439
respiration, 439
chromosome rearrangement,
mtDNA, 837
phylogeography, 837
Cicadidae,
bioacoustics, 815
cryptic species, 815
Tibicen, 815
Cicindela limbata labradorensis,
Cicindela limbata nogahabarensis, 277
mitochondrial DNA, 277
subspecies, 277
tiger beetle, 277
Cicindela limbata nogahabarensis,
Cicindela limbata labradorensis, 277
mitochondrial DNA, 277
subspecies, 277
tiger beetle, 277
Cimicomorpha,
Curalium, 20
Heteroptera, 20
new genus, 20
phylogenetics, 20
circadian rhythm,
clock gene, 1121
melon fly, 1121
quality control, 1121
sterile insect techniques, 1121
classification,
Coenaletoidea, 501
Collembola, 501
Entomobryomorpha, 501
new superfamily, 501
climate,
biological control, 539
phenology, 539
Solenopsis invicta, 539
species relationships, 539
clock gene,
circadian rhythm, 1121
melon fly, 1121
quality control, 1121
sterile insect techniques, 1121
cloud forests,
biodiversity, 128
conservation, 128
Curculionidae, 128
tropical montane forests, 128
Clusia,
Amuroclusia syn. nov., 327
Clusiidae, 327
Melanoclusia gen. nov., 327
Paraclusia syn. nov., 327
Clusiidae,
Amuroclusia syn. nov., 327
Clusia, 327
Melanoclusia gen. nov., 327
Paraclusia syn. nov., 327
coastal animals,
Coleoptera, 939
phylogeographic breaks, 939
serial founder event, 939
Tenebrionidae, 939
codling moth,
apple, 675
attraction, 675
host-finding, 675
pear, 675
Coenaletoidea,
classification, 501
Collembola, 501
Entomobryomorpha, 501
new superfamily, 501
coevolution,
fungal disease, 113
parasitism, 113
predation, 113
Spiroplasma kunkelii, 113
COI, 491, 845
28S-D2, 491
COII, 845
Dioryctria, 845
identification, 491
mitochondrial DNA, 845
Neohydatothrips, 491

phylogeny, 491
seed orchard, 845
COII,
COI, 845
Dioryctria, 845
mitochondrial DNA, 845
seed orchard, 845
COII gene,
cuticular hydrocarbons, 763
range extension, 763
COII barcoding,
Africa paleobiogeography, 978
cryptic speciation, 978
interspecific hybridization, 978
wing shape, 978
cold-free space,
Coleomegilla maculata lengi, 860
Harmonia axyridis, 860
overwintering survival, 860
Coleomegilla maculata lengi,
cold-free space, 860
Harmonia axyridis, 860
overwintering survival, 860
Coleoptera, 53, 402, 411, 474, 939
abundance, 402
Atlantic forest, 402
Bolboceratini, 474
coastal animals, 939
conservation, 402
dune conservation, 411
Geotrupidae, 474
key, 53
new species, 53
off-road vehicle, 411
Orizabus, 53
phylogeographic breaks, 939
Pseudocotalpa, 411
richness, 402
Scarabaeidae, 53
Scarabaeoidea, 474
serial founder event, 939
Tenebrionidae, 939
collective dispersal,
amplification process, 247
solitary behavior, 247
Tetranychus urticae, 247
trail following, 247
Collembola,
classification, 501
Coenaletoidea, 501
Entomobryomorpha, 501
new superfamily, 501
columnar epithelial cells,
histology, 159
insect midgut, 159
light microscopy, 159
morphology, 159
community gardens,
insect survey, 140
invasive species, 140
pollinator conservation, 140
urban ecology, 140
competition strategy,
aggression, 449
endophagous insect, 449
intraspecific competition, 449
pupal cell, 449
complete mitochondrial DNA sequence,
Archaeognatha, 1131
inversion, 1131
Pedetontus silvestrii, 1131
conjugated linoleic acid,
fatty acids, 430
Ostrinia nubilalis, 430
Conopomorpha,
cytochrome oxidase I, 930
elongation factor-1α, 930
genetic bottleneck, 930
Theobroma cacao, 930
Conotrachelus nenuphar,
mtCOI, 824
phylogeography, 824
polymorphism, 824
strains, 824
conservation, 128, 402, 833, 1078
Abies guatemalensis, 833
abundance, 402
agricultural, 1078
aphid, 833
aphidophagous, 1078
Atlantic forest, 402
biodiversity, 128
biological control, 1078
cloud forests, 128
Coleoptera, 402
Curculionidae, 128
Guatemala, 833
invasive, 1078
richness, 402
tropical montane forests, 128
conspecific sperm precedence,
Allonemobius, 30
ground cricket, 30
hybrid zone, 30
hybridization, 30
contact chemoreceptors,
antennae, 1094
electron microscopy, 1094
sensilla, 1094
Coptotermes formosanus,
amino acid, 1088
feeding deterrent, 1088
wood preservative, 1088
Corethrella appendiculata,
Aedes albopictus, 1150
predator–prey, 1150
size-dependent, 1150
courtship,
behavioral sequences, 215
harlequin bug, 215
Hemiptera, 215
Pentatomidae, 215
courtship song,
C. carnea group, 971
cytochrome oxidase subunit I, 971
diagnostic marker, 971
mitochondrial polyphyly, 971
cricket,
calling song, 260
mate choice, 260
predation, 260
Zapotitlán Salinas, 260
cryptic speciation,
Africa paleobiogeography, 978
COII barcoding, 978
interspecific hybridization, 978
wing shape, 978
cryptic species, 743, 815
agriculturally important species, 743
bioacoustics, 815
Cicadidae, 815
mitochondrial DNA, 743
morphology, 743
phylogeny, 743
Tibicen, 815
Culex territans,
amphibian, 95
attraction, 95
mosquito, 95
mosquito hearing, 95
Curalium,
Cimicomorpha, 20
Heteroptera, 20
new genus, 20
phylogenetics, 20
Curculionidae, 7, 128
biodiversity, 128
cloud forests, 128
conservation, 128
new species, 7
taxonomy, 7
tropical montane forests, 128
Turkey, 7
yellow starthistle, 7
cuticular hydrocarbons, 598, 763
aphid epicuticule, 598
COII gene, 763
exuviae, 598
range extension, 763
Sitobion avenae, 598
wax esters, 598
cytochrome oxidase I, 525, 619, 930
Conopomorpha, 930
elongation factor-1α, 525, 930
fall armyworm, 619
genetic bottleneck, 930
mitochondrial DNA, 619
Planococcus, 525
polymerase chain reaction-restriction fragment

length polymorphism, 619
Pseudococcidae, 525
taxonomy, 525
Theobroma cacao, 930
cytochrome oxidase subunit I,
C. carnea group, 971
courtship song, 971
diagnostic marker, 971
mitochondrial polyphyly, 971

D

degeneration,
midgut epithelium, 151
necrosis, 151
regeneration, 151
degree-days,
biological control, 604
Blissidae, 604
developmental rate, 604
Poaceae, 604
Delphastus catalinae,
Bemisia tabaci, 378
biological control, 378
predator, 378
survival, 378
Dendroctonus murrayanae,
lodgepole pine, 1010
Scolytinae, 1010
density dependence,
host range, 70
mutualism, 70
Publilia modesta, 70
spatial variation, 70
development,
life table, 1017
Liposcelis decolor, 1017
temperature, 1017
developmental rate,
biological control, 604
Blissidae, 604
degree-days, 604
Poaceae, 604
developmental stage,
Dyar's rule, 881
instars, 881
southwestern Andalusia, 881
Streblote panda, 881
diagnostic marker,
C. carnea group, 971
courtship song, 971
cytochrome oxidase subunit I, 971
mitochondrial polyphyly, 971
diapause,
chilling, 439
Melanoplus, 439
respiration, 439
diapause induction,
Microplitis mediator, 613
photoperiod, 613
temperature, 613
Dimocarpomyia folicola,
Asphonyliini, 301
Dimocarpus longan, 301
gall midges, 301
Thailand, 301
Dimocarpus longan,
Asphonyliini, 301
Dimocarpomyia folicola, 301
gall midges, 301
Thailand, 301
Dioryctria,
COI, 845
COII, 845
mitochondrial DNA, 845
seed orchard, 845
dispersal, 197, 899
AFLP, 197
Cactoblastis cactorum, 899
gene flow, 197
Grapholita molesta, 197
introduction, 899
invasive species, 899
phylogeography, 899
population genetic structure, 197
distribution map,
Blattella asahinai, 397
new distribution records, 397
peridomestic pest, 397
β diversity,
habitat specificity, 362
parasitoid diversity, 362
species richness host associations, 362
Tachinidae, 362
DNA molecular marker,
Ceratitis capitata, 627
introduction sources, 627
mitochondrial haplotypes, 627
population dynamics, 627
Dromiciocoptes,
external morphology, 289
mammalian parasites, 289
myocoptidae, 289
systematics, 289
drone,
bees, 1137
brood, 1137
hygiene, 1137
varroa, 1137
droplet production,
blow fly, 964
bubbling, 964
insect crop, 964
regurgitation, 964
dumping behavior,
Callosobruchus maculatus, 950
egg-laying decision, 950
host range, 950
dune conservation,
Coleoptera, 411
off-road vehicle, 411
Pseudocotalpa, 411
dung beetles,
assemblages, 58
southern alps, 58
spatial and temporal segregation, 58
Dyar's rule,
developmental stage, 881
Dyar's rule, 881
instars, 881
southwestern Andalusia, 881
Streblote panda, 881

E

Early Carboniferous,
Australia, 79
orthopteroid, 79
pteridosperm, 79
time lag, 79
ecology,
Caribbean, 1026
Erythraeidae, 1026
Opiliones, 1026
parasitism, 1026
egg case,
life history, 689
morphology, 689
predation, 689
egg-laying decision,
Callosobruchus maculatus, 950
dumping behavior, 950
host range, 950
electron microscopy, 1094, 1103
antennae, 1094
contact chemoreceptors, 1094
sensilla, 1094
sensilla basiconica, 1103
sensilla chaetica, 1103
sensory fields, 1103
uniporous sensilla, 1103
elongation factor-1α, 525, 930
Conopomorpha, 930
cytochrome oxidase I, 525, 930
genetic bottleneck, 930
Planococcus, 525
Pseudococcidae, 525
taxonomy, 525
Theobroma cacao, 930
endophagous insect,
aggression, 449
competition strategy, 449
intraspecific competition, 449
pupal cell, 449
Entomobryomorpha,
classification, 501
Coenaletoidea, 501
Collembola, 501
new superfamily, 501
enzyme-linked immunosorbent assay,
gut contents, 887
insect predation, 887

monoclonal antibodies, 887
Erythraeidae,
Caribbean, 1026
ecology, 1026
Opiliones, 1026
parasitism, 1026
Eublemminae,
Araeopteroninae, 40
Glenoides lenticuligera, 40
Hypenodinae, 40
Nigetia formosalis, 40
Eueides,
Callophrys, 786
Heliconius, 786
sexual coevolution, 786
spermatophore, 786
excavating behavior,
tunnel architecture, 456
wood consumption, 456
external morphology,
Dromiciocoptes, 289
mammalian parasites, 289
myocoptidae, 289
systematics, 289
extirpated species,
ant collection techniques, 1056
faunal survey, 1056
pest species, 1056
state list, 1056
exuviae,
aphid epicuticule, 598
cuticular hydrocarbons, 598
Sitobion avenae, 598
wax esters, 598

F

facultative diapause,
monophagous herbivore, 392
phenotypic plasticity, 392
seasonal change in host quality, 392
fall armyworm, 619, 639
cytochrome oxidase I, 619
isolation, 639
mitochondrial DNA, 619
polymerase chain reaction-restriction fragment length polymorphism, 619
reproductive compatibility, 639
strain, 639
fatty acids,
conjugated linoleic acid, 430
Ostrinia nubilalis, 430
faunal survey,
ant collection techniques, 1056
extirpated species, 1056
pest species, 1056
state list, 1056
feeding deterrent,
amino acid, 1088
Coptotermes formosanus, 1088
wood preservative, 1088
Figitidae,
Hymenoptera, 993
Neralsia, 993
new species, 993
North America, 993
finite rate of increase,
net reproductive rate, 1033
pollen as food, 1033
spider mites as prey, 1033
thrips as prey, 1033
flightless Lepidoptera,
bagworm, 872
population regulation, 872
Thyridopteryx ephemeraeformis, 872
white pine, 872
food,
Calosoma sayi, 104
Carabidae, 104
Spodoptera frugiperda, 104
starvation, 104
foraging,
activity, 272
foraging effort, 272
foraging success, 272
Pogonomyrmex occidentalis, 272
foraging effort,
activity, 272
foraging, 272
foraging success, 272
Pogonomyrmex occidentalis, 272
foraging success,
activity, 272
foraging, 272
foraging effort, 272
Pogonomyrmex occidentalis, 272
forest landscapes,
ant communities, 119
imported fire ants, 119
null models, 119
Formica,
homing, 1145
orientation, 1145
polarized light, 1145
Frankliniella spp.,
population dynamics, 769
thrips, 769
fruit borer,
Annona cherimola, 181
antennal sensilla, 181
Talponia batesi, 181
fruit fly,
Bactrocera invadens, 331
host plants, 331
host preference, 331
infestation index, 331
fungal disease,
coevolution, 113
parasitism, 113
predation, 113
Spiroplasma kunkelii, 113

G

Galendromus (*Mugidromus*),
Brazil, 804
carinulatus species group, 804
Galendromus (*Mugidromus*), 804
predatory mites, 804
South America, 804
gall midges, 297, 301
Asphonyliini, 301
Dimocarpomyia folicola, 301
Dimocarpus longan, 301
Obolodiplosis robiniae, 297
Platygaster robiniae, 297
platygastridae, 297
Robinia pseudoacacia, 297
Thailand, 301
gene flow,
AFLP, 197
dispersal, 197
Grapholita molesta, 197
population genetic structure, 197
genetic bottleneck,
Conopomorpha, 930
cytochrome oxidase I, 930
elongation factor-1α, 930
Theobroma cacao, 930
genitalia,
introgression, 794
phylogeography, 794
sibling species, 794
Sierra Nevada, 794
Geotrupidae,
Bolboceratini, 474
Coleoptera, 474
Scarabaeoidea, 474
gibberellic acid,
Apanteles galleriae, 593
parasitoid, 593
risk assessment, 593
toxicity, 593
Glenoides lenticuligera,
Araeopteroninae, 40
Eublemminae, 40
Hypenodinae, 40
Nigetia formosalis, 40
Gramineae,
Heteroptera, 558
host plant, 558
host range, 558
rice bug, 558
Grapholita molesta,
AFLP, 197
dispersal, 197

gene flow, 197
population genetic structure, 197
grasshoppers,
adult survival, 922
B-chromosomes, 922
body-size related traits, 922
gray sunflower seed weevil,
nested clade analysis, 204
polymorphic microsatellite loci, 204
16S rDNA, 204
greenhouses,
biological control, 384
modified Malaise trap, 384
population dynamics, 384
Syrphidae, 384
ground cricket,
Allonemobius, 30
conspecific sperm precedence, 30
hybrid zone, 30
hybridization, 30
Guatemala,
Abies guatemalensis, 833
aphid, 833
conservation, 833
gut contents,
enzyme-linked immunosorbent assay, 887
insect predation, 887
monoclonal antibodies, 887
gut endosymbiont,
Alcaligenes, 1050
blood-feeding moth, 1050
Klebsiella, 1050
microbial associate, 1050

H

habitat specificity,
β diversity, 362
parasitoid diversity, 362
species richness host associations, 362
Tachinidae, 362
haplotype,
polymorphisms, 1112
Spodoptera frugiperda, 1112
harlequin bug,
behavioral sequences, 215
courtship, 215
Hemiptera, 215
Pentatomidae, 215
Harmonia axyridis,
cold-free space, 860
Coleomegilla maculata lengi, 860
overwintering survival, 860
Heliconius,
Callophrys, 786
Eueides, 786
sexual coevolution, 786
spermatophore, 786
Helotidae,
new species, 722
taxonomic revision, 722
vigorsii species group, 722
Hemiptera, 215, 307
behavioral sequences, 215
courtship, 215
harlequin bug, 215
heteroptera, 307
lygaeoidea, 307
Mexico, 307
Pentatomidae, 215
Xestocoris, 307
Heteroptera, 20, 307, 558
Cimicomorpha, 20
Curalium, 20
Gramineae, 558
hemiptera, 307
host plant, 558
host range, 558
lygaeoidea, 307
Mexico, 307
new genus, 20
phylogenetics, 20
rice bug, 558
Xestocoris, 307
histology,
columnar epithelial cells, 159
insect midgut, 159
light microscopy, 159
morphology, 159
Homalodisca coagulata,
acoustic signal, 253
leafhopper, 253
sharpshooter, 253
substrate vibration, 253
homing,
Formica, 1145
orientation, 1145
polarized light, 1145
host discrimination,
learning, 656
oviposition behavior, 656
Tephritidae, 656
host plant, 331, 558, 906
ant-organs, 906
Bactrocera invadens, 331
balloon setae, 906
fruit fly, 331
Gramineae, 558
Heteroptera, 558
host preference, 331
host range, 558
infestation index, 331
myrmecoxeny, 906
rice bug, 558
scanning electron microscopy, 906
host preference,
Bactrocera invadens, 331
fruit fly, 331
host plants, 331
infestation index, 331
host range, 70, 558, 950
Callosobruchus maculatus, 950
density dependence, 70
dumping behavior, 950
egg-laying decision, 950
Gramineae, 558
Heteroptera, 558
host plant, 558
mutualism, 70
Publilia modesta, 70
rice bug, 558
spatial variation, 70
host selection,
Nicrophorus investigator, 266
phoresy, 266
phoretic mites, 266
host selection behavior,
chemical ecology, 955
chemical stimuli, 955
physical stimuli, 955
host size model,
Musca domestica, 650
parasitoid, 650
sex ratio, 650
host-finding,
apple, 675
attraction, 675
codling moth, 675
pear, 675
huanglongbing,
Asian citrus psyllid, 585
body color, 585
pigment, 585
reproductive state, 585
hybrid zone,
Allonemobius, 30
conspecific sperm precedence, 30
ground cricket, 30
hybridization, 30
hybridization, 30, 172
Allonemobius, 30
conspecific sperm precedence, 30
ground cricket, 30
hybrid zone, 30
Lycaeides, 172
morphology, 172
Nabokov, 172
Remington suture zones, 172
hygiene,
bees, 1137
brood, 1137
drone, 1137
varroa, 1137

Hymenoptera,
Figitidae, 993
Neralsia, 993
new species, 993
North America, 993
Hypenodinae,
Araeopteroninae, 40
Eublemminae, 40
Glenoides lenticuligera, 40
Nigetia formosalis, 40

I

identification,
COI, 491
Neohydatothrips, 491
phylogeny, 491
28S-D2, 491
identification key,
Neanuridae, 683
Paranura, 683
taxonomy, 683
immature stages,
biological control, 189
Braconidae, 189
larval development, 189
parasitoid, 189
imported fire ants,
ant communities, 119
forest landscapes, 119
null models, 119
infestation index,
Bactrocera invadens, 331
fruit fly, 331
host plants, 331
host preference, 331
insect,
bees, 915
midgut, 915
peritrophic membrane, 915
insect crop,
blow fly, 964
bubbling, 964
droplet production, 964
regurgitation, 964
insect development,
insect phenology, 351
nonlinear development model, 351
Russian wheat aphid, 351
same-shape development distribution, 351
insect midgut,
columnar epithelial cells, 159
histology, 159
light microscopy, 159
morphology, 159
insect phenology,
insect development, 351
nonlinear development model, 351
Russian wheat aphid, 351
same-shape development distribution, 351
insect predation,
enzyme-linked immunosorbent assay, 887
gut contents, 887
monoclonal antibodies, 887
insect survey,
community gardens, 140
invasive species, 140
pollinator conservation, 140
urban ecology, 140
Insecta,
bioecology, 779
cassava root scale, 779
Manihot esculenta, 779
instars,
developmental stage, 881
Dyar's rule, 881
southwestern Andalusia, 881
Streblote panda, 881
interference competition,
ants, 1162
Linepithema, 1162
Solenopsis, 1162
venom, 1162
interspecific hybridization,
Africa paleobiogeography, 978
COII barcoding, 978
cryptic speciation, 978
wing shape, 978
intraspecific competition,
aggression, 449
competition strategy, 449
endophagous insect, 449
pupal cell, 449
intraspecific variation,
polymerase chain reaction, 565
introduction,
Cactoblastis cactorum, 899
dispersal, 899
invasive species, 899
phylogeography, 899
introduction sources,
Ceratitis capitata, 627
DNA molecular marker, 627
mitochondrial haplotypes, 627
population dynamics, 627
introgression,
genitalia, 794
phylogeography, 794
sibling species, 794
Sierra Nevada, 794
invasive,
agricultural, 1078
aphidophagous, 1078
biological control, 1078
conservation, 1078
invasive species, 140, 341, 899
Cactoblastis cactorum, 899
community gardens, 140
dispersal, 899
insect survey, 140
introduction, 899
phylogeography, 899
pollinator conservation, 140
predators, 341
soybean aphid, 341
urban ecology, 140
inversion,
Archaeognatha, 1131
complete mitochondrial DNA sequence, 1131
Pedetontus silvestrii, 1131
isolation,
fall armyworm, 639
reproductive compatibility, 639
strain, 639
ITS1,
ITS2, 514
phylogeography, 514
Sphaeropthalminae, 514
ITS2,
ITS1, 514
phylogeography, 514
Sphaeropthalminae, 514

K

key,
Coleoptera, 53
new species, 53
Orizabus, 53
Scarabaeidae, 53
Klebsiella,
Alcaligenes, 1050
blood-feeding moth, 1050
gut endosymbiont, 1050
microbial associate, 1050

L

larvae,
larval behavior, 1041
larval morphology, 1041
pupae, 1041
sap feeders, 1041
larval behavior,
larvae, 1041
larval morphology, 1041
pupae, 1041
sap feeders, 1041
larval development,
biological control, 189
Braconidae, 189
immature stages, 189
parasitoid, 189
larval morphology, 13, 1041
larvae, 1041
larval behavior, 1041
Pselaphinae, 13

pupae, 1041
rove beetles, 13
sap feeders, 1041
short-winged mold beetles, 13
Staphylinidae, 13
leafhopper, 253, 1003
acoustic signal, 253
Homalodisca coagulata, 253
overwintering strategy, 1003
Rosa chinensis, 1003
Rubus gr. *fruticosus*, 1003
sharpshooter, 253
substrate vibration, 253
Vitis vinifera, 1003
learning,
host discrimination, 656
oviposition behavior, 656
Tephritidae, 656
Leptinotarsa decemlineata,
sperm competition, 371
sperm precedence, 371
sterile male technique, 371
Leucochrysini,
Mata Atlântica, 314
new genus, 314
new species, 314
Santocellus atlanticis, 314
life cycle,
caddisfly, 573
case building, 573
phenology, 573
protandry, 573
life history,
egg case, 689
morphology, 689
predation, 689
life table,
development, 1017
Liposcelis decolor, 1017
temperature, 1017
light,
mate guarding, 1154
mating behavior, 1154
sperm competition, 1154
temperature, 1154
light microscopy,
columnar epithelial cells, 159
histology, 159
insect midgut, 159
morphology, 159
Linepithema,
ants, 1162
interference competition, 1162
Solenopsis, 1162
venom, 1162
Liposcelis decolor,
development, 1017
life table, 1017
temperature, 1017
lodgepole pine,
Dendroctonus murrayanae, 1010
Scolytinae, 1010
Lycaeides,
hybridization, 172
morphology, 172
Nabokov, 172
Remington suture zones, 172
lygaeoidea,
hemiptera, 307
heteroptera, 307
Mexico, 307
Xestocoris, 307

M

macropterous,
alary polymorphism, 551
below-ground insects, 551
brachypterous, 551
male genitalia,
androconial system, 465
caddisfly, 465
Neotropics, 465
scales, 465
mammalian parasites,
Dromiciocoptes, 289
external morphology, 289
myocoptidae, 289
systematics, 289
Manihot esculenta,
bioecology, 779
cassava root scale, 779
Insecta, 779
Mantispidae,
Mexico, 703
Neuroptera, 703
species list, 703
Mata Atlântica,
Leucochrysini, 314
new genus, 314
new species, 314
Santocellus atlanticis, 314
mate choice,
calling song, 260
cricket, 260
predation, 260
Zapotitlán Salinas, 260
mate guarding,
light, 1154
mating behavior, 1154
sperm competition, 1154
temperature, 1154
mating behavior, 1154, 1169
light, 1154
mate guarding, 1154
Podisus maculiventris, 1169
song repertoire, 1169
sperm competition, 1154
substrate-borne communication, 1169
temperature, 1154
mating competitiveness,
Bactrocera tryoni, 664
sterile insect technique, 664
Megaloptera,
Neotropics, 808
Protosialis, 808
Sialidae, 808
taxonomy, 808
Melanoclusia gen. nov.,
Amuroclusia syn. nov., 327
Clusia, 327
Clusiidae, 327
Paraclusia syn. nov., 327
Melanoplus,
chilling, 439
diapause, 439
respiration, 439
melon fly,
circadian rhythm, 1121
clock gene, 1121
quality control, 1121
sterile insect techniques, 1121
Mexico, 307, 703
hemiptera, 307
heteroptera, 307
lygaeoidea, 307
Mantispidae, 703
Neuroptera, 703
species list, 703
Xestocoris, 307
microbial associate,
Alcaligenes, 1050
blood-feeding moth, 1050
gut endosymbiont, 1050
Klebsiella, 1050
Microplitis mediator,
diapause induction, 613
photoperiod, 613
temperature, 613
midgut,
bees, 915
insect, 915
peritrophic membrane, 915
midgut epithelium,
degeneration, 151
necrosis, 151
regeneration, 151
mitochondrial DNA, 277, 619, 743, 845
agriculturally important species, 743
Cicindela limbata labradorensis, 277
Cicindela limbata nogahabarensis, 277
COI, 845
COII, 845
cryptic species, 743
cytochrome oxidase I, 619
Dioryctria, 845
fall armyworm, 619
morphology, 743
phylogeny, 743
polymerase chain reaction-restriction fragment

length polymorphism, 619
seed orchard, 845
subspecies, 277
tiger beetle, 277
mitochondrial haplotypes,
Ceratitis capitata, 627
DNA molecular marker, 627
introduction sources, 627
population dynamics, 627
mitochondrial polyphyly,
C. carnea group, 971
courtship song, 971
cytochrome oxidase subunit I, 971
diagnostic marker, 971
modified Malaise trap,
biological control, 384
greenhouses, 384
population dynamics, 384
Syrphidae, 384
molecular systematics,
Apidae, 713
Neotropical, 713
parasitoid, 713
monoclonal antibodies,
enzyme-linked immunosorbent assay, 887
gut contents, 887
insect predation, 887
monophagous herbivore,
facultative diapause, 392
phenotypic plasticity, 392
seasonal change in host quality, 392
morphology, 159, 172, 689, 743
agriculturally important species, 743
columnar epithelial cells, 159
cryptic species, 743
egg case, 689
histology, 159
hybridization, 172
insect midgut, 159
life history, 689
light microscopy, 159
Lycaeides, 172
mitochondrial DNA, 743
Nabokov, 172
phylogeny, 743
predation, 689
Remington suture zones, 172
mosquito,
amphibian, 95
attraction, 95
Culex territans, 95
mosquito hearing, 95
mosquito hearing,
amphibian, 95
attraction, 95
Culex territans, 95
mosquito, 95
mtCOI,
Conotrachelus nenuphar, 824
phylogeography, 824
polymorphism, 824
strains, 824
mtDNA,
chromosome rearrangement, 837
phylogeography, 837
Musca domestica,
host size model, 650
parasitoid, 650
sex ratio, 650
mutualism,
density dependence, 70
host range, 70
Publilia modesta, 70
spatial variation, 70
myocoptidae,
Dromiciocoptes, 289
external morphology, 289
mammalian parasites, 289
systematics, 289
myrmecoxeny,
ant-organs, 906
balloon setae, 906
host plants, 906
scanning electron microscopy, 906

N

Nabokov,
hybridization, 172
Lycaeides, 172
morphology, 172
Remington suture zones, 172
Neanuridae,
identification key, 683
Paranura, 683
taxonomy, 683
necrosis,
degeneration, 151
midgut epithelium, 151
regeneration, 151
Neohydatothrips,
COI, 491
identification, 491
phylogeny, 491
28S-D2, 491
Neotropical,
Apidae, 713
molecular systematics, 713
parasitoid, 713
Neotropics, 465, 808
androconial system, 465
caddisfly, 465
male genitalia, 465
Megaloptera, 808
Protosialis, 808
scales, 465
Sialidae, 808
taxonomy, 808
Neralsia,
Figitidae, 993
Hymenoptera, 993
new species, 993
North America, 993
nested clade analysis,
gray sunflower seed weevil, 204
polymorphic microsatellite loci, 204
16S rDNA, 204
net reproductive rate,
finite rate of increase, 1033
pollen as food, 1033
spider mites as prey, 1033
thrips as prey, 1033
Neuroptera,
Mantispidae, 703
Mexico, 703
species list, 703
new distribution records,
Blattella asahinai, 397
distribution map, 397
peridomestic pest, 397
new genus, 20, 314
Cimicomorpha, 20
Curalium, 20
Heteroptera, 20
Leucochrysini, 314
Mata Atlântica, 314
new species, 314
phylogenetics, 20
Santocellus atlanticis, 314
new species, 7, 53, 314, 722, 993
Coleoptera, 53
Curculionidae, 7
Figitidae, 993
Helotidae, 722
Hymenoptera, 993
key, 53
Leucochrysini, 314
Mata Atlântica, 314
Neralsia, 993
new genus, 314
North America, 993
Orizabus, 53
Santocellus atlanticis, 314
Scarabaeidae, 53
taxonomic revision, 722
taxonomy, 7
Turkey, 7
vigorsii species group, 722
yellow starthistle, 7
new superfamily,
classification, 501
Coenaletoidea, 501
Collembola, 501
Entomobryomorpha, 501
Nicrophorus investigator,
host selection, 266
phoresy, 266

phoretic mites, 266
Nigetia formosalis,
Araeopteroninae, 40
Eublemminae, 40
Glenoides lenticuligera, 40
Hypenodinae, 40
nonlinear development model,
insect development, 351
insect phenology, 351
Russian wheat aphid, 351
same-shape development distribution, 351
North America,
Figitidae, 993
Hymenoptera, 993
Neralsia, 993
new species, 993
null models,
ant communities, 119
forest landscapes, 119
imported fire ants, 119

O

O,O,O-trimethyl phosphorothioate,
O,O,S-trimethyl phosphorothioate, 418
O,S,S-trimethyl phosphorodithioate, 418
volatile malathion, 418
O,O,S-trimethyl phosphorothioate,
O,O,O-trimethyl phosphorothioate, 418
O,S,S-trimethyl phosphorodithioate, 418
volatile malathion, 418
O,S,S-trimethyl phosphorodithioate,
O,O,O-trimethyl phosphorothioate, 418
O,O,S-trimethyl phosphorothioate, 418
volatile malathion, 418
Obolodiplosis robiniae,
gall midges, 297
Platygaster robiniae, 297
platygastridae, 297
Robinia pseudoacacia, 297
off-road vehicle,
Coleoptera, 411
dune conservation, 411
Pseudocotalpa, 411
Opiliones,
Caribbean, 1026
ecology, 1026
Erythraeidae, 1026
parasitism, 1026
orientation,
Formica, 1145
homing, 1145
polarized light, 1145
Orizabus,
Coleoptera, 53
key, 53
new species, 53
Scarabaeidae, 53
orthopteroid,
Australia, 79
Early Carboniferous, 79
pteridosperm, 79
time lag, 79
Ostrinia nubilalis,
conjugated linoleic acid, 430
fatty acids, 430
overwintering strategy,
leafhoppers, 1003
Rosa chinensis, 1003
Rubus gr. *fruticosus*, 1003
Vitis vinifera, 1003
overwintering survival,
cold-free space, 860
Coleomegilla maculata lengi, 860
Harmonia axyridis, 860
oviposition behavior,
host discrimination, 656
learning, 656
Tephritidae, 656

P

Paraclusia syn. nov.,
Amuroclusia syn. nov., 327
Clusia, 327
Clusiidae, 327
Melanoclusia gen. nov., 327
Paranura,
identification key, 683
Neanuridae, 683
taxonomy, 683
parasitic bees,
bee conservation, 1067
biodiversity, 1067
soil-nesting species, 1067
suburban development, 1067
parasitism, 113, 533, 1026
Caribbean, 1026
coevolution, 113
ecology, 1026
Erythraeidae, 1026
fungal disease, 113
Opiliones, 1026
predation, 113
prediction of development, 533
Spiroplasma kunkelii, 113
synchrony of host and parasitoid, 533
temperature, 533
thermal constants for development, 533
parasitoid, 189, 593, 650, 713
Apanteles galleriae, 593
Apidae, 713
biological control, 189
Braconidae, 189
gibberellic acid, 593
host size model, 650
immature stages, 189
larval development, 189
molecular systematics, 713
Musca domestica, 650
Neotropical, 713
risk assessment, 593
sex ratio, 650
toxicity, 593
parasitoid diversity,
β diversity, 362
habitat specificity, 362
species richness host associations, 362
Tachinidae, 362
parasitoid wasp,
remating, 229
Spalangia endius, 229
unattractiveness, 229
virgin females, 229
pear,
apple, 675
attraction, 675
codling moth, 675
host-finding, 675
Pedetontus silvestrii,
Archaeognatha, 1131
complete mitochondrial DNA sequence, 1131
inversion, 1131
Pentatomidae, 215, 235
acoustic communication, 235
behavioral sequences, 215
courtship, 215
harlequin bug, 215
Hemiptera, 215
sex-specific signal, 235
substrate-borne vibrational signal, 235
peridomestic pest,
Blattella asahinai, 397
distribution map, 397
new distribution records, 397
peritrophic membrane,
bees, 915
insect, 915
midgut, 915
pest species,
ant collection techniques, 1056
extirpated species, 1056
faunal survey, 1056
state list, 1056
phenology, 539, 573
biological control, 539
caddisfly, 573
case building, 573
climate, 539
life cycle, 573

protandry, 573
Solenopsis invicta, 539
species relationships, 539
phenotypic plasticity,
facultative diapause, 392
monophagous herbivore, 392
seasonal change in host quality, 392
phoresy,
host selection, 266
Nicrophorus investigator, 266
phoretic mites, 266
phoretic mites,
host selection, 266
Nicrophorus investigator, 266
phoresy, 266
photoperiod,
diapause induction, 613
Microplitis mediator, 613
temperature, 613
phylogenetics,
Cimicomorpha, 20
Curalium, 20
Heteroptera, 20
new genus, 20
phylogeny, 491, 743
agriculturally important species, 743
COI, 491
cryptic species, 743
identification, 491
mitochondrial DNA, 743
morphology, 743
Neohydatothrips, 491
28S-D2, 491
phylogeographic breaks,
coastal animals, 939
Coleoptera, 939
serial founder event, 939
Tenebrionidae, 939
phylogeography, 514, 794, 824, 837, 899
Cactoblastis cactorum, 899
chromosome rearrangement, 837
Conotrachelus nenuphar, 824
dispersal, 899
genitalia, 794
introduction, 899
introgression, 794
invasive species, 899
ITS1, 514
ITS2, 514
mtCOI, 824
mtDNA, 837
polymorphism, 824
sibling species, 794
Sierra Nevada, 794
Sphaeropthalminae, 514
strains, 824
physical stimuli,
chemical ecology, 955
chemical stimuli, 955
host selection behavior, 955
pigment,
Asian citrus psyllid, 585
body color, 585
huanglongbing, 585
reproductive state, 585
Planococcus,
cytochrome oxidase I, 525
elongation factor 1α, 525
Pseudococcidae, 525
taxonomy, 525
Platygaster robiniae,
gall midges, 297
Obolodiplosis robiniae, 297
platygastridae, 297
Robinia pseudoacacia, 297
platygastridae,
gall midges, 297
Obolodiplosis robiniae, 297
Platygaster robiniae, 297
Robinia pseudoacacia, 297
Poaceae,
biological control, 604
Blissidae, 604
degree-days, 604
developmental rate, 604
Podisus maculiventris,
mating behavior, 1169
song repertoire, 1169
substrate-borne communication, 1169
Pogonomyrmex occidentalis,
activity, 272
foraging, 272
foraging effort, 272
foraging success, 272
polarized light,
Formica, 1145
homing, 1145
orientation, 1145
pollen as food,
finite rate of increase, 1033
net reproductive rate, 1033
spider mites as prey, 1033
thrips as prey, 1033
pollinator conservation,
community gardens, 140
insect survey, 140
invasive species, 140
urban ecology, 140
polymerase chain reaction,
intraspecific variation, 565
polymerase chain reaction-restriction fragment length polymorphism,
cytochrome oxidase I, 619
fall armyworm, 619
mitochondrial DNA, 619
polymorphic microsatellite loci,
gray sunflower seed weevil, 204
nested clade analysis, 204
16S rDNA, 204
polymorphism, 824, 1112
Conotrachelus nenuphar, 824
haplotype, 1112
mtCOI, 824
phylogeography, 824
Spodoptera frugiperda, 1112
strains, 824
population dynamics, 384, 627, 769
biological control, 384
Ceratitis capitata, 627
DNA molecular marker, 627
Frankliniella spp., 769
greenhouses, 384
introduction sources, 627
mitochondrial haplotypes, 627
modified Malaise trap, 384
Syrphidae, 384
thrips, 769
population genetic structure,
AFLP, 197
dispersal, 197
gene flow, 197
Grapholita molesta, 197
population regulation,
bagworm, 872
flightless Lepidoptera, 872
Thyridopteryx ephemeraeformis, 872
white pine, 872
predation, 113, 260, 689
calling song, 260
coevolution, 113
cricket, 260
egg case, 689
fungal disease, 113
life history, 689
mate choice, 260
morphology, 689
parasitism, 113
Spiroplasma kunkelii, 113
Zapotitlán Salinas, 260
predator, 341, 378
Bemisia tabaci, 378
biological control, 378
Delphastus catalinae, 378
invasive species, 341
soybean aphid, 341
survival, 378
predator-prey,
Aedes albopictus, 1150
Corethrella appendiculata, 1150
predator-prey, 1150
size-dependent, 1150
predatory mites,
Brazil, 804
carinulatus species group, 804

Galendromus (*Mugidromus*), 804
South America, 804
prediction of development,
parasitism, 533
synchrony of host and parasitoid, 533
temperature, 533
thermal constants for development, 533
protandry,
caddisfly, 573
case building, 573
life cycle, 573
phenology, 573
Protosialis,
Megaloptera, 808
Neotropics, 808
Sialidae, 808
taxonomy, 808
Pselaphinae,
larval morphology, 13
rove beetles, 13
short-winged mold beetles, 13
Staphylinidae, 13
Pseudococcidae,
cytochrome oxidase I, 525
elongation factor 1α, 525
Planococcus, 525
taxonomy, 525
Pseudocotalpa,
Coleoptera, 411
dune conservation, 411
off-road vehicle, 411
pteridosperm,
Australia, 79
Early Carboniferous, 79
orthopteroid, 79
time lag, 79
Publilia modesta,
density dependence, 70
host range, 70
mutualism, 70
spatial variation, 70
pupae,
larvae, 1041
larval behavior, 1041
larval morphology, 1041
sap feeders, 1041
pupal cell,
aggression, 449
competition strategy, 449
endophagous insect, 449
intraspecific competition, 449

Q

quality control,
circadian rhythm, 1121
clock gene, 1121
melon fly, 1121
sterile insect techniques, 1121

R

range extension,
COII gene, 763
cuticular hydrocarbons, 763
recurrent event,
Apis mellifera, 867
survival, 867
swarming, 867
Varroa destructor, 867
regeneration,
degeneration, 151
midgut epithelium, 151
necrosis, 151
regurgitation,
blow fly, 964
bubbling, 964
droplet production, 964
insect crop, 964
remating,
parasitoid wasp, 229
Spalangia endius, 229
unattractiveness, 229
virgin females, 229
Remington suture zones,
hybridization, 172
Lycaeides, 172
morphology, 172
Nabokov, 172
reproductive compatibility,
fall armyworm, 639
isolation, 639
strain, 639
reproductive state,
Asian citrus psyllid, 585
body color, 585
huanglongbing, 585
pigment, 585
respiration,
chilling, 439
diapause, 439
Melanoplus, 439
rice bug,
Gramineae, 558
Heteroptera, 558
host plant, 558
host range, 558
richness,
abundance, 402
Atlantic forest, 402
Coleoptera, 402
conservation, 402
risk assessment,
Apanteles galleriae, 593
gibberellic acid, 593
parasitoid, 593
toxicity, 593
Robinia pseudoacacia,
gall midges, 297
Obolodiplosis robiniae, 297
Platygaster robiniae, 297
platygastridae, 297
Rosa chinensis,
leafhoppers, 1003
overwintering strategy, 1003
Rubus gr. *fruticosus*, 1003
Vitis vinifera, 1003
rove beetles,
larval morphology, 13
Pselaphinae, 13
short-winged mold beetles, 13
Staphylinidae, 13
Rubus gr. *fruticosus*,
leafhoppers, 1003
overwintering strategy, 1003
Rosa chinensis, 1003
Vitis vinifera, 1003
Russian wheat aphid,
insect development, 351
insect phenology, 351
nonlinear development model, 351
same-shape development distribution, 351

S

16S rDNA,
gray sunflower seed weevil, 204
nested clade analysis, 204
polymorphic microsatellite loci, 204
28S-D2,
COI, 491
identification, 491
Neohydatothrips, 491
phylogeny, 491
same-shape development distribution,
insect development, 351
insect phenology, 351
nonlinear development model, 351
Russian wheat aphid, 351
Santocellus atlanticis,
Leucochrysini, 314
Mata Atlântica, 314
new genus, 314
new species, 314
sap feeders,
larvae, 1041
larval behavior, 1041
larval morphology, 1041
pupae, 1041
SC,
springtails, 989
taxonomy, 989
scales,
androconial system, 465
caddisfly, 465
male genitalia, 465
Neotropics, 465

scanning electron microscopy,
ant-organs, 906
balloon setae, 906
host plants, 906
myrmecoxeny, 906
Scarabaeidae,
Coleoptera, 53
key, 53
new species, 53
Orizabus, 53
Scarabaeoidea,
Bolboceratini, 474
Coleoptera, 474
Geotrupidae, 474
Scolytinae,
Dendroctonus murrayanae, 1010
lodgepole pine, 1010
seasonal change in host quality,
facultative diapause, 392
monophagous herbivore, 392
phenotypic plasticity, 392
seed orchard,
COI, 845
COII, 845
Dioryctria, 845
mitochondrial DNA, 845
sensilla,
antennae, 1094
contact chemoreceptors, 1094
electron microscopy, 1094
sensilla basiconica,
electron microscopy, 1103
sensilla chaetica, 1103
sensory fields, 1103
uniporous sensilla, 1103
sensilla chaetica,
electron microscopy, 1103
sensilla basiconica, 1103
sensory fields, 1103
uniporous sensilla, 1103
sensory fields,
electron microscopy, 1103
sensilla basiconica, 1103
sensilla chaetica, 1103
uniporous sensilla, 1103
serial founder event,
coastal animals, 939
Coleoptera, 939
phylogeographic breaks, 939
Tenebrionidae, 939
sex ratio,
host size model, 650
Musca domestica, 650
parasitoid, 650
sex-specific signal,
acoustic communication, 235
Pentatomidae, 235
substrate-borne vibrational signal, 235
sexual coevolution,
Callophrys, 786
Eueides, 786
Heliconius, 786
spermatophore, 786
sharpshooter,
acoustic signal, 253
Homalodisca coagulata, 253
leafhopper, 253
substrate vibration, 253
short-winged mold beetles,
larval morphology, 13
Pselaphinae, 13
rove beetles, 13
Staphylinidae, 13
Sialidae,
Megaloptera, 808
Neotropics, 808
Protosialis, 808
taxonomy, 808
sibling species,
genitalia, 794
introgression, 794
phylogeography, 794
Sierra Nevada, 794
Sierra Nevada,
genitalia, 794
introgression, 794
phylogeography, 794
sibling species, 794
Sitobion avenae,
aphid epicuticule, 598
cuticular hydrocarbons, 598
exuviae, 598
wax esters, 598
size-dependent,
Aedes albopictus, 1150
Corethrella appendiculata, 1150
predator–prey, 1150
soil-nesting species,
bee conservation, 1067
biodiversity, 1067
parasitic bees, 1067
suburban development, 1067
Solenopsis,
ants, 1162
interference competition, 1162
Linepithema, 1162
venom, 1162
Solenopsis invicta,
biological control, 539
climate, 539
phenology, 539
species relationships, 539
solitary behavior,
amplification process, 247
collective dispersal, 247
Tetranychus urticae, 247
trail following, 247
song repertoire,
mating behavior, 1169
Podisus maculiventris, 1169
substrate-borne communication, 1169
South America,
Brazil, 804
carinulatus species group, 804
Galendromus (*Mugidromus*), 804
predatory mites, 804
southern alps,
assemblages, 58
dung beetles, 58
spatial and temporal segregation, 58
southwestern Andalusia,
developmental stage, 881
Dyar's rule, 881
instars, 881
Streblote panda, 881
soybean aphid,
invasive species, 341
predators, 341
Spalangia endius,
parasitoid wasp, 229
remating, 229
unattractiveness, 229
virgin females, 229
spatial and temporal segregation,
assemblages, 58
dung beetles, 58
southern alps, 58
spatial variation,
density dependence, 70
host range, 70
mutualism, 70
Publilia modesta, 70
species list,
Mantispidae, 703
Mexico, 703
Neuroptera, 703
species relationships,
biological control, 539
climate, 539
phenology, 539
Solenopsis invicta, 539
species richness host associations,
β diversity, 362
habitat specificity, 362
parasitoid diversity, 362
Tachinidae, 362
sperm competition, 371, 1154
Leptinotarsa decemlineata, 371
light, 1154
mate guarding, 1154
mating behavior, 1154
sperm precedence, 371
sterile male technique, 371
temperature, 1154
sperm precedence,
Leptinotarsa decemlineata, 371
sperm competition, 371
sterile male technique, 371

spermatophore,
Callophrys, 786
Eueides, 786
Heliconius, 786
sexual coevolution, 786
Sphaeropthalminae,
ITS1, 514
ITS2, 514
phylogeography, 514
spider mites as prey,
finite rate of increase, 1033
net reproductive rate, 1033
pollen as food, 1033
thrips as prey, 1033
Spiroplasma kunkelii,
coevolution, 113
fungal disease, 113
parasitism, 113
predation, 113
Spodoptera frugiperda, 104, 1112
Calosoma sayi, 104
Carabidae, 104
food, 104
haplotype, 1112
polymorphisms, 1112
starvation, 104
springtails,
SC, 989
taxonomy, 989
Staphylinidae,
larval morphology, 13
Pselaphinae, 13
rove beetles, 13
short-winged mold beetles, 13
starvation,
Calosoma sayi, 104
Carabidae, 104
food, 104
Spodoptera frugiperda, 104
state list,
ant collection techniques, 1056
extirpated species, 1056
faunal survey, 1056
pest species, 1056
sterile insect technique, 664, 1121
Bactrocera tryoni, 664
circadian rhythm, 1121
clock gene, 1121
mating competitiveness, 664
melon fly, 1121
quality control, 1121
sterile male technique,
Leptinotarsa decemlineata, 371
sperm competition, 371
sperm precedence, 371
strain, 639, 824
Conotrachelus nenuphar, 824
fall armyworm, 639
isolation, 639
mtCOI, 824
phylogeography, 824
polymorphism, 824
reproductive compatibility, 639
Streblote panda,
developmental stage, 881
Dyar's rule, 881
instars, 881
southwestern Andalusia, 881
subspecies,
Cicindela limbata labradorensis, 277
Cicindela limbata nogahabarensis, 277
mitochondrial DNA, 277
tiger beetle, 277
substrate vibration,
acoustic signal, 253
Homalodisca coagulata, 253
leafhopper, 253
sharpshooter, 253
substrate-borne communication,
mating behavior, 1169
Podisus maculiventris, 1169
song repertoire, 1169
substrate-borne vibrational signal,
acoustic communication, 235
Pentatomidae, 235
sex-specific signal, 235
suburban development,
bee conservation, 1067
biodiversity, 1067
parasitic bees, 1067
soil-nesting species, 1067
survival, 378, 867
Apis mellifera, 867
Bemisia tabaci, 378
biological control, 378
Delphastus catalinae, 378
predator, 378
recurrent event, 867
swarming, 867
Varroa destructor, 867
swarming,
Apis mellifera, 867
recurrent event, 867
survival, 867
Varroa destructor, 867
synchrony of host and parasitoid,
parasitism, 533
prediction of development, 533
temperature, 533
thermal constants for development, 533
Syrphidae,
biological control, 384
greenhouses, 384
modified Malaise trap, 384
population dynamics, 384
systematics,
Dromiciocoptes, 289
external morphology, 289
mammalian parasites, 289
myocoptidae, 289

T

Tachinidae,
β diversity, 362
habitat specificity, 362
parasitoid diversity, 362
species richness host associations, 362
Talponia batesi,
Annona cherimola, 181
antennal sensilla, 181
fruit borer, 181
taxonomic revision,
Helotidae, 722
new species, 722
vigorsii species group, 722
taxonomy, 7, 525, 683, 808, 989
Curculionidae, 7
cytochrome oxidase I, 525
elongation factor 1α, 525
identification key, 683
Megaloptera, 808
Neanuridae, 683
Neotropics, 808
new species, 7
Paranura, 683
Planococcus, 525
Protosialis, 808
Pseudococcidae, 525
SC, 989
Sialidae, 808
springtails, 989
Turkey, 7
yellow starthistle, 7
temperature, 533, 613, 1017, 1154
development, 1017
diapause induction, 613
life table, 1017
light, 1154
Liposcelis decolor, 1017
mate guarding, 1154
mating behavior, 1154
Microplitis mediator, 613
parasitism, 533
photoperiod, 613
prediction of development, 533
sperm competition, 1154
synchrony of host and parasitoid, 533
thermal constants for development, 533
Tenebrionidae,
coastal animals, 939
Coleoptera, 939
phylogeographic breaks, 939
serial founder event, 939
Tephritidae,
host discrimination, 656
learning, 656

oviposition behavior, 656
Tetranychus urticae,
amplification process, 247
collective dispersal, 247
solitary behavior, 247
trail following, 247
Thailand,
Asphonyliini, 301
Dimocarpomyia folicola, 301
Dimocarpus longan, 301
gall midges, 301
Theobroma cacao,
Conopomorpha, 930
cytochrome oxidase I, 930
elongation factor-1α, 930
genetic bottleneck, 930
thermal constants for development,
parasitism, 533
prediction of development, 533
synchrony of host and parasitoid, 533
temperature, 533
thrips,
Frankliniella spp., 769
population dynamics, 769
thrips as prey,
finite rate of increase, 1033
net reproductive rate, 1033
pollen as food, 1033
spider mites as prey, 1033
Thyridopteryx ephemeraeformis,
bagworm, 872
flightless Lepidoptera, 872
population regulation, 872
white pine, 872
Tibicen,
bioacoustics, 815
Cicadidae, 815
cryptic species, 815
tiger beetle,
Cicindela limbata labradorensis, 277
Cicindela limbata nogahabarensis, 277
mitochondrial DNA, 277
subspecies, 277
time lag,
Australia, 79
Early Carboniferous, 79
orthopteroid, 79
pteridosperm, 79
toxicity,
Apanteles galleriae, 593
gibberellic acid, 593
parasitoid, 593
risk assessment, 593
trail following,
amplification process, 247
collective dispersal, 247
solitary behavior, 247
Tetranychus urticae, 247
tropical montane forests,
biodiversity, 128
cloud forests, 128
conservation, 128
Curculionidae, 128
tunnel architecture,
excavating behavior, 456
wood consumption, 456
Turkey,
Curculionidae, 7
new species, 7
taxonomy, 7
yellow starthistle, 7

U

unattractiveness,
parasitoid wasp, 229
remating, 229
Spalangia endius, 229
virgin females, 229
uniporous sensilla,
electron microscopy, 1103
sensilla basiconica, 1103
sensilla chaetica, 1103
sensory fields, 1103
urban ecology,
community gardens, 140
insect survey, 140
invasive species, 140
pollinator conservation, 140

V

varroa,
bees, 1137
brood, 1137
drone, 1137
hygiene, 1137
Varroa destructor,
Apis mellifera, 867
recurrent event, 867
survival, 867
swarming, 867
venom,
ants, 1162
interference competition, 1162
Linepithema, 1162
Solenopsis, 1162
vigorsii species group,
Helotidae, 722
new species, 722
taxonomic revision, 722
virgin females,
parasitoid wasp, 229
remating, 229
Spalangia endius, 229
unattractiveness, 229
Vitis vinifera,
leafhoppers, 1003
overwintering strategy, 1003
Rosa chinensis, 1003
Rubus gr. *fruticosus*, 1003
volatile malathion,
O,O,O-trimethyl phosphorothioate, 418
O,O,S-trimethyl phosphorothioate, 418
O,S,S-trimethyl phosphorodithioate, 418

W

wax esters,
aphid epicuticule, 598
cuticular hydrocarbons, 598
exuviae, 598
Sitobion avenae, 598
white pine,
bagworm, 872
flightless Lepidoptera, 872
population regulation, 872
Thyridopteryx ephemeraeformis, 872
wing shape,
Africa paleobiogeography, 978
COII barcoding, 978
cryptic speciation, 978
interspecific hybridization, 978
wood consumption,
excavating behavior, 456
tunnel architecture, 456
wood preservative,
amino acid, 1088
Coptotermes formosanus, 1088
feeding deterrent, 1088

X

Xestocoris,
hemiptera, 307
heteroptera, 307
lygaeoidea, 307
Mexico, 307

Y

yellow starthistle,
Curculionidae, 7
new species, 7
taxonomy, 7
Turkey, 7

Z

Zapotitlán Salinas,
calling song, 260
cricket, 260
mate choice, 260
predation, 260